



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

**Another Richardson's Owl in Massachusetts.**—As *Nyctala tengmalmi richardsoni* has been so seldom taken so far south as this point (central Eastern Massachusetts), it may be well to record an additional example. I have in my collection a fine specimen obtained in this place on Jan. 1, 1885. It was approached without difficulty in broad daylight by a wood-chopper and killed with a stick.

This is about the southern limit of the range of the species as at present determined. I find but three examples on record to the south of this; the early one of Dr. Wood for Connecticut in 1859, and the two obtained in 1881 and 1882 near Providence, R. I., as reported by Mr. Jencks in the 'Nuttall Bulletin' of April, 1881, and April, 1883.

The circumstances of this capture indicate the defective day-vision characteristic of the strictly nocturnal species, in accordance with Richardson's statement: "It is so much dazzled by the light of the sun that it becomes stupid and may easily be caught by hand."—F. C. BROWNE, Framingham, Mass.

**The Oyster-catcher (*Hematopus palliatus*) in Massachusetts.**—Mr. Warren Hapgood tells me that during the last week of April, 1885, he received an Oyster-catcher in the flesh from Chatham, Mass. It was shot on Monomoy Island by Alonzo Nye, the veteran gunner of Chatham. It was in fine plumage, and is now preserved in Mr. Hapgood's collection. If I remember rightly this is only the third specimen known to have been taken in Massachusetts, the other two being the pair mentioned by Dr. Brewer (Water Birds. Vol. I, p. 113), and now in the New England collection of the Boston Society.—WILLIAM BREWSTER, Cambridge, Mass.

**The Baird's Sandpiper (*Actodromas bairdii*) at Locust Grove, New York.**—August 18, 1885, while Dr. C. H. Merriam and the writer were driving along a country road at the above locality we discovered a solitary Baird's Sandpiper on the edge of a small, temporary pool of water formed by the recent rains. It was a female of the year in fine plumage, and was evidently a straggling migrant which had dropped down hap-hazard to feed and rest. Most of the eastern stragglers of this species hitherto recorded have been from the Atlantic coast, but undoubtedly the bird occurs more or less commonly all the way across the interior country.—H. W. HENSHAW, Washington, D. C.

**A Bird New to Massachusetts.**—Among the Sandpipers collected by me years ago in Massachusetts I find a single specimen of the Western Sandpiper (*Ereunetes occidentalis*), taken on Long Island, Boston Harbor, Aug. 27, 1870. It was one of a number of 'Peeps' shot on that day, and it is by no means unlikely that the lot contained others of this species, which were overlooked under the impression that they were all the common species, *E. pusillus*. Like the Baird's Sandpiper, this species may be expected to occur in small numbers along the Atlantic coast during the migrations, especially in fall.—H. W. HENSHAW, Washington, D. C.